

DIABETES MELLITUS

(Adapted from Veterinary Information Network www.vin.com)

What is Diabetes Mellitus?

The cells of the body require a sugar known as glucose for food and they depend on the bloodstream to bring glucose to them. They cannot, however, absorb and utilize glucose without a hormone known as insulin. This hormone, insulin, is produced by the pancreas. Insulin is like a key that unlocks the door to separate cells from the sugars in our bloodstream.

Glucose comes from the diet. When an animal goes without food, the body must break down fat, stored starches, and protein to supply calories for the hungry cells. Proteins and starches may be converted into glucose. Fat, however, requires different processing that can lead to the production of ketones rather than glucose. Ketones are another type of fuel that the body can use in a pinch but the detection of ketones indicates that something wrong is happening in the patient's metabolism. Ketones may therefore be detected in the urine of starving animals because of massive fat mobilization is required for ketone formation. Ketones can also be detected in diabetic ketoacidosis, a severe complication of unregulated diabetes so it is helpful to periodically monitor for ketones in a diabetic patient's urine. The point is, for now, that in times of extreme fat burning (such as in starvation), ketones are a byproduct.

The main clinical signs of diabetes mellitus are:

- Excessive eating
- Excessive drinking
- Excessive urination
- Weight loss

It is usually fairly clear from the history and tests showing dramatic glucose elevations in the blood (and usually glucose in the urine, too) that diabetes mellitus is the diagnosis. Some pets are able to substantially raise their blood sugars from stress (such as might occur when a sensitive, sick, and anxious patient goes the vet's office). This could create misleading test results. If there is any question about the diagnosis, a test called a fructosamine level may be requested. This test reflects an average blood glucose level over the past several weeks so if this is also elevated, a one-time elevated glucose can be distinguished from the persistent elevations of true diabetes mellitus. The fructosamine test is also sometimes used in monitoring therapy for diabetes mellitus.

Another common symptom of diabetes mellitus is urinary tract infection. All the sugar in the urine makes the bladder an excellent incubator for bacteria. Antibiotics are necessary to clear up such an infection and some monitoring may be needed to help detect these infections.

Type I and Type II Diabetes Mellitus

Diabetes mellitus is a classical disease in humans and most of us have heard some of the terms used to describe it. In humans, diabetes is broken down into two forms: Type I and Type II. These are also referred to as juvenile onset and adult onset diabetes, or insulin dependent and non-insulin dependent diabetes. In short, Type 1 is the type where the pancreas produces no insulin at all, and in Type 2 the pancreas produces some insulin but not enough. Virtually all dogs have insulin dependent diabetes and must be treated with insulin. Most cats have non-insulin dependent diabetes.

This might suggest that most cats can get away without insulin injections but that is not the case at all. Instead, for cats, there is potential for the diabetes to actually resolve if the pancreas improves its insulin-secreting ability. Insulin injections are needed to treat most diabetic cats but for some cats, the situation is mild enough for oral medication to suffice. Good glucose control and proper diet can resolve the diabetes in some lucky cats.

What happens once a diagnosis is reached

First, an insulin type and dose will need to be selected. There are several types of insulins and it is not possible to know how much insulin your individual pet will require. Your veterinarian can make a guess based on what works for other cats and dogs and what has been reported in the literature. Most pets require injections twice a day, approximately 12 hours apart, following a meal.

You will need to learn how to give insulin injections to your pet. The technique of subcutaneous (under the skin) insulin administration should be thoroughly demonstrated by your vet. You may be surprised to find that the most common reason for a pet having difficulty achieving regulation is that the owner is not giving the injections properly. Be sure you know how to hold the bottle, manipulate the syringe, hold your pet, and give the injection. Some situations require that the pet be hospitalized for a few days for the initial regulation, but most of the time your pet will be at home receiving injections shortly after the diagnosis has been reached. Injections may seem intimidating at first but most pet owners are surprised at how easy this turns out to be.

Insulin must be purchased from the vet. You will also need syringes and a bottle of insulin to begin home treatment. Insulin syringes are marked in insulin units so the insulin syringes must match the insulin concentrations (either U-100 syringes for 100 unit/cc insulins or U-40 syringes for 40 unit/cc insulins.) Always double check these numbers whenever you receive more supplies.

Never alter the insulin dose recommended by your vet. To determine whether dose adjustments are needed (or if a different type of insulin is more appropriate), your pet will need a glucose curve where blood sugar levels are monitored every 2 to 4 hours or so for 12 to 24 hours. This kind of testing tells the doctor how long the insulin injection is lasting as well as what the lowest and highest glucose levels of the day are. It is important to find out when your pet's curve is due. Often in the beginning, it takes several dose selections and several curves before the right dose is determined.

A bottle of insulin, when stored properly, should last 6 to 8 weeks. After that time it should probably be replaced.



DIABETES MELLITUS

(Adapted from Veterinary Information Network www.vin.com)

Home testing

If your pet is too sensitive for a valid glucose curve at the vet's office, the fructosamine blood test may be particularly useful. Again, this test looks at average glucose levels so wide fluctuations will not be discovered but at least there is a monitoring option for this situation.

Ketostix are used to detect ketones in urine and can be obtained at any drug store. If it is not difficult to access your pet's urine, a first morning test is helpful. Remember, finding ketones occasionally is not a problem but a positive dipstick three days in a row is a criterion for a vet visit.

Monitoring

Your pet will probably require re-regulation at some point. During re-regulation periods, expect a curve to be run a week or two after each adjustment in insulin dose. Bring your pet in for a recheck and glucose curve if your pet:

- seems to feel ill
- is losing weight
- has a ravenous appetite or loses its appetite
- seems to be drinking or urinating excessively
- becomes disoriented or groggy
- has ketones in the urine for three days in a row.

Dental work

It is important for diabetic pets to have their teeth cleaned annually. Dental tartar seeds the body with bacteria and when blood sugar levels run high, infections in important organs can take root. The kidneys and heart are particularly vulnerable.



Insulin Shock

If your pet appears wobbly or drunken, the blood sugar level may have dropped too low. This occurs after an insulin overdose. First try to get your pet to eat. If the pet will not eat, administer 1 tablespoon of honey, or even sugar-water. If no improvement occurs, immediately see your veterinarian for emergency treatment. When your pet is more stable, a glucose curve will be needed to determine why this happened and what a more appropriate insulin dose might be.

Difficult regulation

Your pet will probably require re-regulation at some point. There may be an underlying reason to sort out. If your pet seems to fit in this category, some reasons could be:

Improper administration of insulin. If possible, have your doctor observe you giving the insulin to your pet. Another possibility is that your insulin may be out of date.

Rapid insulin metabolism. Insulin wears off quickly in some animals. Your pet may require a different type of insulin or a second injection during the day or even additional injections during the day.

Insulin overdose may actually lead to elevated glucose levels (and clinical signs of diabetes mellitus). In this situation, too much insulin brings the blood glucose too low and other hormones respond to bring it back up (and generally over-do it).

Steroid administration (such as prednisone, prednisolone, etc.) will interfere with insulin.

Progesterone, a female hormone, also interferes with insulin. Unspayed female diabetics should be spayed once they are sufficiently regulated.

Feeding a diabetic cat

Regulation is achieved via a balance of diet, exercise, and insulin. Realizing that therapeutic diets are not always attractive to pets, there are some ideal foods which should at least be offered.

The most up-to-date choice for cats is a low carbohydrate high protein diet. These diets promote weight loss in obese diabetics and are available in both canned and dry formulations such as Science Diet m/d. Talk to your veterinarian to select an appropriate choice for your pet.

Avoid soft-moist diets as sugars are used as preservatives. Avoid breads and sweet treats. If it is not possible to change the pet's diet, then regulation will just have to be worked out around whatever the pet will eat.